

تولید و تعیین خصوصیت بیوشیمیایی فیتاز گرمادوست جدا شده از باکتری باسیلوس آمیلولیکوفاسینس LOR10

ارسطو بدویی دلفارد و مریم پرهام فر

گروه زیست شناسی، دانشکده علوم، دانشگاه شهید باهنر کرمان، کرمان، ایران

مسئول مکاتبات: ارسطو بدویی دلفارد، badoei@uk.ac.ir

چکیده. فیتاز ارزش تغذیه‌ای غذاهای گیاهی را با افزایش هضم‌پذیری پروتئین و در دسترس‌سازی مواد معدنی از طریق هیدرولیز فیتات در طی هضم در معده یا در طی پردازش غذا بهبود می‌بخشد. منابع میکروبی برای تولید تجاری فیتاز پتانسیل قابل توجهی دارند. بنابراین، هدف این تحقیق غربالگری و جداسازی باکتری-های مولد فیتاز از چشمه آب گرم با پتانسیل قابل توجه می‌باشد. شناسایی بهترین سویه با روش مولکولی 16S rDNA انجام شد. بهینه‌سازی تولید آنزیم در حضور منابع مختلف کربن، نیتروژن و فسفات انجام گردید. فعالیت و پایداری آنزیم در pH ها، دماها و یون‌های مختلف انجام شد. مقایسه توالی ژن 16S rDNA سویه LOR10 با سایر سویه‌ها در بانک ژنی با استفاده از Clustal omega میزان ۹۸ درصد همولوژی با باسیلوس آمیلولیکوفاسینس را نشان داد. نتایج بهینه‌سازی محیط نشان داد که گالاکتوز، عصاره مخمر و تری‌کلسیم فسفات به‌عنوان بهترین منبع کربن، نیتروژن و فسفات برای تولید فیتاز می‌باشند. دمای بهینه فعالیت در دمای ۷۰ درجه سانتی‌گراد به دست آمد. بهترین پایداری فیتاز نیز در pH حدود ۵ تا ۸ به دست آمد. فعالیت فیتازی در حضور کلرید-کلسیم، کلریدروی و سولفات‌منیزیم حدود ۱/۴، ۲/۳ و ۱/۶ برابر به ترتیب افزایش یافت. خاطر نشان می‌شود که فعالیت فیتازی در حضور EDTA و SDS حدود ۳۰ درصد کاهش یافت. این نتایج خاطر نشان می‌سازد که فیتاز LOR10 پتانسیل قابل توجهی برای استفاده‌های تجاری از جمله مکمل غذایی را داراست.

واژه‌های کلیدی. باکتری‌های تجزیه‌کننده فیتات، جداسازی، چشمه آب گرم، شناسایی باکتریایی

Production and biochemical characterization of a thermostable phytase from *Bacillus amyloliquefaciens* LOR10

Arastoo Badoei-Dalfard & Maryam Parhamfar

Department of Biology, Faculty of Sciences, Shahid Bahonar University of Kerman, Kerman, Iran

Correspondent author: Arastoo Badoei-Dalfard, badoei@uk.ac.ir

Abstract. Phytase can improve the nutritional value of plant-based foods by enhancing protein digestibility and mineral availability through phytate digestion in the stomach and the food processing industry. Microbial sources are more promising for the production of phytases on a commercial scale. The objectives of this exploration were to screening and isolation of phytase-producing bacteria from hot spring with commercial interest. Molecular identification of the best isolate was achieved by the 16S rDNA gene. Optimization of phytase production was prepared in the presence of different phosphate, nitrogen, and carbon sources. Enzyme activity and stability were also explored in the presence of different pHs, temperatures, and ion compounds. Comparing the 16S rDNA gene sequence of the isolate LOR10 with those in GenBank using Clustal omega shows 98% sequence homology with *Bacillus amyloliquefaciens*. Medium optimization studies showed that galactose, yeast extract, and tricalcium phosphate were the best sources of carbon, nitrogen, and phosphate for phytase production, respectively. The optimum temperature activity was also observed to be 70 °C. Phytase stability was at its optimum in a pH range of 5.0–8.0. Phytase activity increased in the presence of CaCl₂, ZnCl₂, and MnSO₄ about 1.4, 2.3 and 1.6 folds, respectively. It could be mentioned that phytase activity decreased by about 30 % in the presence of EDTA and SDS. On the basis of the results, it could be concluded that LOR10 phytase has a great potential for commercial interest as an additive to animal plant-based foods.

Keywords. bacterial identification, hot spring, isolation, phytate-degrading bacteria

INTRODUCTION

Phytase catalyzes the hydrolysis of phytate (myo-inositol-1,2,3,4,5,6-hexakis dihydrogen phosphate) through phytate digestion in the stomach and the food processing industry (In et al., 2004; Sapna & Singh, 2016). Most of the plant seeds and their co-products contain 1–2% phytic acid representing >60% of their total phosphorus (Lei & Porres, 2003; Puppala, et al., 2019). In monogastric or agastric animals, phytic acid is a polyanionic chelating agent that forms complexes with several divalent cations of primary nutritional importance, e.g., Ca^{2+} , Mg^{2+} , Zn^{2+} , Cu^{2+} , Fe^{2+} , and Mn^{2+} (Gessler et al., 2018). Phytic acid can also form complexes with proteins and amino acids at both acidic and alkaline pHs (Saani et al., 2019). The resulting compounds are insoluble, difficult for humans to hydrolyze during digestion, and, thus are typically nutritionally less available for absorption (Afinah et al., 2010). Furthermore, the unutilized phytate from plant feeds becomes an environmental pollutant in the areas of intensive animal agriculture (Vasudevan et al., 2017). These problems can be overcome by supplementing the animal feed with phytases, which improve the nutritional status of the feed and further, declines environmental phosphorus pollution caused by the use of phosphorus in animal feed (Selle & Ravindran, 2008; Singha & Satyanarayana, 2011; Demir et al., 2018; Vasudevan et al., 2019).

Phosphate can be efficiently released from the phytate by phytase activity (Kumar et al., 2010). Phytase blended into the diets of poultry, swine, and fish to increase the availability of phosphorus, minerals, amino acids, and energy. The enzyme sequesters orthophosphate groups from the inositol ring of phytate to produce free inorganic phosphate (Baruah et al., 2007; Cao et al., 2007). Phytases are present in bacteria, fungi, animals, and plants. Commonly, the activity of an animal's phytase is insignificant compared to the phytase of microorganisms and plants (Lei et al., 2013). Phytases have been found in various microorganisms such as *Aspergillus fumigatus* (Vasudevan et al., 2017), *Escherichia coli* (Demir et al., 2018), *Enterobacter* sp.4 (Vasudevan et al., 2019), *B. amyloliquefaciens* (Selle & Ravindran, 2008), *Lactobacillus sanfranciscensis* (Raghavendra & Halami, 2009), *Bacillus* sp. (Choi et al., 2001), *Raoultella* sp. (Sajidan et al., 2004), *Citrobacter braakii* (Kim et al., 1998), and *Mitsuokella jalaludinii* (Lan et al., 2002). Owing to some properties such as catalytic productivity, substrate specificity, and resistance to proteolytic enzymes, bacterial phytases are a real alternative to the fungal enzymes (Konietzny & Greiner, 2004).

Because of their great industrial significance, there is an ongoing interest in isolating new microbial producing phytase and optimization of enzyme production. Hence the aim of this study is the isolation of the bacteria with phytase producing ability and characterization of this enzyme.

MATERIALS AND METHODS

Isolation of the phytate-degrading bacteria

Samples were gathered from the Loriya hot spring in Jiroft, which located in the southeast of Iran. Approximately 5ml of these samples was injected in 50 ml of the phytase-specific medium [1.5% glucose, 0.5% $(\text{NH}_4)_2\text{SO}_4$, 0.05% KCl, 0.01% $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01% NaCl, 0.01% $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.001% FeSO_4 , 0.001% MnSO_4 , pH 6.5 with 0.5% sodium phytate (Sigma)]. These flasks were maintained at 45 °C, with 180 rpm for three days. After that, 5ml of this culture was added to the same fresh medium and maintained in the same conditions. Finally, 0.1 ml of this culture was streaked onto the phytase-specific medium and incubated at 65 °C for 48h. Bacterial colonies capable of hydrolyzing sodium phytate, which can be recognized by their surrounding clear halo, were obtained by re-plating single colonies (Chunshan et al., 2001). To find the optimum time of phytase production, *Bacillus* sp. LOR10 was cultured in phytase specific medium, samples were picked up at 24 h intervals, and the phytase activity was measured as described below.

Measurement of phytase activity

The phytase activity was investigated using a modified ferrous sulfate molybdenum blue system (Greiner et al., 1997). Briefly, 50 μl of the enzyme solution was incubated with 450 μl of a substrate solution (1.5 mM sodium phytate in 0.25 M sodium acetate buffer, pH 5.0) at 45 °C for 30 min. The reaction was then stopped by adding 0.5 ml of 10% (w/v) trichloroacetic acid. After that, the released inorganic phosphate was analyzed by adding 1ml of color reagent, containing 1% (w/v) ammonium molybdate, 3.2% (v/v) sulfuric acid, and 7.2% (w/v) ferrous sulfate, and the optical density measured at 700 nm. All the phytase activity determinations were accomplished in three independent experiments. One unit (U) of phytase activity was defined as the amount of enzyme that released 1 micromole of phosphate per minute at 45 °C. The investigational error was not ever over 7%.

PCR amplification and 16S rDNA sequencing

Genomic DNA of *Bacillus* sp. LOR10 was extracted according to Sambrook and Russell protocols (Sambrook & Russell, 2001), and its purity was checked by the A260/A280. Universal 16S rDNA PCR forward primer 27F (5-

AGTTTGATCCTGGCTCAG-3) and reverse primer 1492R (5-GGC/TACCTTGTTACGACTT-3) were used for the amplification of *16S rDNA* genes (Badoei-Dalfard & Karami, 2013; Ramezani-pour et al., 2015). PCR program was achieved as follows: (1) 94 °C for 5 min as initial temperature, (2) a run of 30 cycles with each cycle consisting of 45 s at 94 °C, 45 s at 48 °C and 90 s at 72 °C, and (3) 5 min at 72 °C to permit for the extension of any incomplete products (Azadian et al., 2016; Farahmand et al., 2019). PCR products were electrophoresed on agarose gel (1.0 %) and subsequently amplified *16S rDNA* bands were purified by DNA extraction kit (Sinaclon, Iran). Then DNA sequencing was performed on both strands directly by SEQ-LAB (Germany).

The phylogenetic tree was prepared based on the comparison of *16S rDNA* sequences of *Bacillus* sp. LOR10 strain with *16S rDNA* sequences of strains of *Bacillus* species stored in the GenBank database (<http://www.ncbi.nlm.nih.gov>). All sequences were aligned with Clustal Omega (version 6.0) that was obtained from <http://www.seqtool.sdsc.edu/CGI/Omega.cgi> (Sievers et al., 2011), and the phylogenetic tree was prepared in MEGA5 (Tamura et al., 2007). The obtained *16S rDNA* sequence has been deposited in GenBank for isolate *Bacillus* sp. LOR10.

Examination of culture condition for phytase production

To determine the optimal condition for phytase production, *Bacillus* sp. LOR10 was inoculated into 50 ml of the liquid medium in a 250 ml Erlenmeyer flask and maintained on a shaker at 37 °C for three days. Carbon, nitrogen, and phosphate sources were explored on the phytase production (Parhamfar et al., 2015; Parhamfar et al., 2016). Carbon sources were 1.5% glucose, 1.5% galactose, and 0.75% glucose + 0.75% galactose. Nitrogen sources were 0.5% glycine, 0.5% yeast extract, and 0.5% ammonium sulfate. Phosphate sources were sodium phytate (0.5%), tricalcium phosphate (0.5%), sodium phytate (0.25%) + tricalcium phosphate (0.25%), and nutrient broth. The effect of initial pH on phytase production was also explored by using phytase specific medium with pH 5.0, 6.0, and 7.0. Samples were picked up each 24h, and phytase activity was determined as described above.

Partial purification of phytase

Bacillus sp. LOR10 was grown up in nutrient broth medium for 16 h with 168 rpm. The liquid medium used for the production of phytase was composed of [1.5% glucose, 0.5% (NH₄)₂SO₄, 0.05% KCl, 0.01% MgSO₄.7H₂O, 0.01% NaCl, 0.01% CaCl₂.2H₂O, 0.001% FeSO₄, 0.001% MnSO₄, pH 6.5 with 0.5% sodium phytate]. The pH of the medium was adjusted to 6.8 before autoclave.

The medium was inoculated at 10% (v/v) with a 16h old culture and incubated at 37 °C with 190 rpm shaking for 48 h. After that, the culture medium was centrifuged at 10,000 × g for 10 min at 4 °C. and subsequently, the supernatant was precipitated with 0–100% (NH₄)₂SO₄ to fraction the proteins at 4 °C. After centrifuging at 12,000×g for 20 min at 4 °C, the pellet was dissolved in a small amount of 10 mM sodium acetate buffer, pH 7.0, and dialyzed overnight against the same buffer. All the steps were carried out at 4 °C. The concentrated sample was applied on a Q-sepharose column (1.5×24 cm), which equilibrated with sodium acetate buffer (20 mM, pH 7.0). The proteins were eluted at a flow rate of 1.0 ml/min with the same buffer with NaCl gradient (0.0-1.0 M). All the steps were carried out at 4 °C. Fractions with high phytase activity were pooled and concentrated. SDS-PAGE was also performed on slab gel containing 10% (w/v) polyacrylamide.

Enzyme characterization

The temperature activity profile of LOR10 phytase was investigated from 30 to 90 °C with 10 °C intervals. The thermal stability profile was also considered by measuring the residual enzyme activity after incubating the LOR10 phytase in 0.25M sodium acetate (pH 6.5) at each temperature for 30 min, and then the enzyme samples placed on ice. Finally, the residual activity was measured as the standard condition.

The pH activity profile was determined by measuring the LOR10 phytase activity at 45 °C, and pH 3.0-11.0 using the following buffers; sodium acetate (0.25M, pH 3.0-6.0), Tris-HCl (0.25M, pH 6.0-9.0), and glycine-NaOH (0.25M, pH 9.0-11). For the pH stability assays, LOR10 phytase was incubated in the same buffers over a range of pH from 3.0 to 11.0 at 45 °C for 60 min. The residual phytase activity was measured under the standard conditions (pH 6.8, 45 °C, 30 min).

LOR10 phytase activity in the presence of different ions was also investigated with the final concentration of 5 mM. For determining ion stability, the equal volumes of metal ions/SDS/EDTA (10 mM) were mixed with enzyme solution in 0.05 M sodium acetate buffer. The mixture was incubated for 1.0 h at room temperature, and then the residual activity was measured as the standard condition.

RESULTS

Isolation and identification of the phytase-producing bacteria

A total of 85 bacterial species were isolated from Loriya hot spring in Jiroft, located in the southeast of Iran. This hot spring has 68 °C temperature and

pH 5.0. After phytase assay screening, the best isolate was selected based on a clear halo around the colony. This isolate was cultured in liquid media, and phytase production was considered. This strain was grown on the liquid medium containing sodium phytate as an inducer. The samples were picked up to measure the phytase activity at a 24h interval. Results showed that the maximum enzyme production occurred after 40 h of incubation (Fig. 1A). It also results in Figure 1B shows that pH

decreased from 6.8 to 5.7 as enzyme production and phosphate solubilization increased.

PCR product of the *16S rDNA* gene was about 1500 bp, and its sequence was edited to a total length of 1260 bp. Comparing the *16S rDNA* gene sequence of the isolate LOR10 with those in GenBank using Clustal omega show 98% sequence homology with *B. amyloliquefaciens* and the phylogenetic tree was created by the neighbor-joining method by MEGA5 software (Fig. 2).




Figure 1. A. Time course of LOR10 phytase production. B. Time course of pH changes during incubation of *Bacillus amyloliquefaciens* LOR10.




Figure 2. *16S rDNA* based phylogenetic analysis of *B. amyloliquefaciens* LOR10. Bootstrap values and scale bar depicting substitution rate per site are indicated. The phylogenetic tree constructed by the neighbor-joining method showing the position of isolate LOR10. *Paenibacillus* sp. 172 was used as an outgroup.

Culture investigation for phytase production

To conclude the best conditions for phytase production, the LOR10 isolate was cultured into 50 ml of the liquid medium in 500 ml flask and maintained on a reciprocal at 37 °C for three days. Figures 3 and 4 displayed diverse sources of carbon, nitrogen, phosphate, and initial pH on phytase production. For carbon sources Glc (1.5%), Gal (1.5%), and Glc (0.75%) + Gal (0.75%) was added to the medium, and phytase activity was assayed at 24, 48, and 72 h of incubation. Results showed that all three carbon sources enhanced phytase production in 24 and 48 h of incubation and marginally decreased in 72h. High phytase activity was gotten in Gal (1.5%) medium after 48h of incubation (Fig. 3a). Results showed that glucose has less effect on phytase production.

The best nitrogen source for phytase production was yeast extract at 48h of incubation. It was also 40 % and 50% more efficient than glycine and ammonium sulfate, respectively (Fig. 3b). Results

showed that glycine had less effect on phytase production among these compounds.

The effect of initial pH on the phytase production by LOR10 isolate was also investigated. Results showed that pH 7.0 was the optimum initial pH for phytase production after 24 h of incubation. But, after 48 and 72 h of incubation, pH 6.0 showed the best results for phytase activity. It is mentioned that after 72h incubation, pH 6 showed 10 and 20% more phytase activity than pH 5.0 and 7.0, respectively (Fig. 4A).

The effect of phosphate salts, such as sodium phytate (0.5%), tricalcium phosphate (0.5%), sodium phytate (0.25%) +tricalcium phosphate (0.25%), and nutrient broth on phytase production is shown on Figure 4b. Results showed that tricalcium phosphate was the most suitable compound for phytase production. The nutrient broth has no significant effect on phytase production. A 30 % enhancement in the phytase production achieved after the addition of tricalcium phosphate to the liquid medium.




Figure 3. Effect of different carbon (A), nitrogen (B) on phytase production of *B. amyloliquefaciens* LOR10. Samples were picked up at 24h intervals and phosphate solubilizing determined as described in material and methods.




Figure 4. Effect of different initial pH (A) and phosphate sources (B) on phytase production of *B. amyloliquefaciens* LOR10. Samples were picked up at 24h intervals and phosphate solubilizing determined as described in material and methods.




Figure 5. SDS-page of purified phytase from *B. amyloliquefaciens* LOR10 (2), protein markers (1).




Figure 6. Temperature activity (A) and stability (B) profiles of LOR10 phytase. For temperature activity, the enzyme was assayed at different temperatures. For temperature stability, the enzyme was pre-incubated at different temperatures for 2h, and then the remaining activity was measured at the standard assay condition, which discussed in the materials and methods.




Figure 7. pH activity (A) and stability (B) profiles of LOR10 phytase. For pH activity, the enzyme was assayed at different pHs. For pH stability, the enzyme was pre-incubated at different pHs for 2h, and then the remaining activity was measured at the standard assay condition, which discussed in the materials and methods.

Table 1. Relative activity and stability of LOR10 phytase in the presence of different ions and compounds

	Relative activity (%)	Relative stability (%)
No ion	100±0	100±0
CaCl ₂	142±1.2	137±1.3
MgCl ₂	105±1.03	112±1.3
ZnCl ₂	232±1.3	125±1.1
KCl	100±1.0	100±1.1
NaCl	100±0.9	152±1.16
MnSO ₄	163±1.4	192±1.5
FeSO ₄	70±0.2	75±0.3
EDTA	65±0.3	90±0.8
SDS	70±0.4	
	60±0.7	

Phytase characterization

SDS-Page results showed that the purified phytase from *B. amyloliquefaciens* LOR10 has a single band of about 22 kDa (Fig. 5). A variety of molecular weight for phytases from other *Bacillus* species had been described as 41 kDa from *Bacillus subtilis* (Farhat et al., 2008), 47 kDa from *Bacillus licheniformes* (Tye et al., 2002), and 28 kDa from *Bacillus tequilensis* Dm018 (Badoei-dalfard et al., 2019).

Phytase activity and stability at different temperatures

The temperature activity profile of LOR10 phytase was shown in Figure 6A. The optimal temperature activity was also detected in 70 °C. LOR10 phytase showed about 35 % activity in 90 °C. Results showed that the phytase activity was constant from 30 to 50 °C, and then regularly decreased (Fig. 6B).

Phytase activity and stability at different pH

The pH activity profile of LOR10 phytase was shown in Figure 7. Results indicated that the LOR10 phytase showed maximum activity at pH 7.0 (Fig. 7A). Also, pH stability results were presented in Figure 7B. These results indicated that LOR10 phytase was stable from pH 5.0–8.0.

Effect of metal ion on phytase activity and stability

LOR10 phytase activity was also investigated at different ions. Results in Table 1 showed that the phytase activity had no change in the presence of MgCl₂, KCl, and NaCl. Phytase activity in the presence of CaCl₂, ZnCl₂, and MnSO₄ was increased about 1.4, 2.3, and 1.6 folds, respectively. It is mentioned that the phytase activity decreased by about 70 % in the presence of EDTA and SDS. Results of ion stability in Table 1 revealed that the phytase activity was improved by about 1.37, 1.52, and 1.92 folds in the presence of CaCl₂, NaCl, and MnSO₄, respectively.

DISCUSSION

A phytase-producing bacterium has been isolated from Loriya hot spring, that it is closely related to

Bacillus amyloliquefaciens. The ideal enzyme creation was revealed after 40 h of cultivation in the liquid medium. Carbon sources investigation results showed that the high phytase activity (100 %) was obtained in galactose medium after two days of incubation, compared to phytase activity (60%) of the glucose medium.

The effect of different parameters on phytase production was also investigated based on previous reports (Parhamfar et al., 2016; Kim et al., 2002). However, phytase production by *Bacillus subtilis* using different carbon sources showed that glucose was the best carbon source (Kim et al., 2002). Nitrogen sources investigation showed that yeast extract had the best effect on the phytase production after 48 h of incubation. Ammonium sulfate and glycine have less effect about 60 and 40 % on phytase production in the same condition. It was also earlier described that the addition of yeast extract (0.01%) to culture improved phosphate solubilizing capacity of *Pseudomonas* sp.2 about 44%. The same results were also reported by Kim and coworkers. They indicated that yeast extract was the better nitrogen source for the phytase production by strain BPTK4 (Kim et al., 2002). The value of phytase production in the presence of different pHs showed that pH 6.0 was more effective than pH 5.0, 7.0 after 72 h of incubation. Interestingly, phytase production was gradually decreased in pH 7.0 but gradually increased in pH 5.0. It was also stated that the optimal pH by *Pseudomonas* sp. YH40 for phytase production was pH 6.0 (Kim et al., 2002). Phosphate sources exploration showed that tricalcium phosphate was the best phosphate source for phytase production. It was previously reported that none of the phosphate salts (calcium phytate, sodium phytate, K₂HPO₄, and Na₂HPO₄) had any effect on increasing phytase production by *Pseudomonas* sp. YH40 (Kim et al., 2002). The addition of KCl and FeSO₄ to the medium showed the same results as *Bacillus* sp.C43 phytase (Sreedevi & Reddy, 2012). LOR10 phytase

has more than 90% of its stability in 40 to 60 °C. This phytase displayed about 60% of its initial activity in 80 °C. The thermal stability of phytase is deliberated to be an important and valuable principle for application as an animal additive. Power and Jagannathan described a phytase that displayed optimal activity at a temperature of 60 °C and was stable up to 70 °C (Powar & Jagannathan, 1982). *Bacillus* sp. (natto) phytase had an optimum temperature of 60 °C (Shimizu, 1992). Although the maximum activity of LOR10 phytase was observed in pH 7.0, it showed more than 40% of its activity in acidic pH (4.0-7.0). LOR10 phytase showed about 90 % of its initial activity in the broad ranges of pH (5.0-8.0). The optimum pH activity of *B. subtilis* (natto) phytase was between 6.0-6.5 (Shimizu, 1992). pH stability of *Pseudomonas* sp. phytase was between pH 5.0-7.0, and the optimal pH of *Pseudomonas* sp. phytase was found pH 7.0 (Kim et al., 2002).

For the investigation of the effects of the ions on phytase activity and stability, several different metal ions were examined (Table 1). Results showed that LOR10 phytase was inhibited by EDTA and SDS about 30%, and stimulated by CaCl₂ (1.4 fold), ZnCl₂ (2.3 fold), and MnSO₄ (1.6 fold). LOR10 phytase stability showed that CaCl₂, ZnCl₂, NaCl, and MnSO₄ stimulated phytase stability from 1.25 to 1.95 folds. It was also reported that Ca²⁺ was required for the activity of *B. subtilis* phytase, and had no critical effect on *Bacillus* phytase (Sreedevi & Reddy, 2012), and had a slightly inhibitory effect on phytases from *E. coli* (Greiner & Konietzny, 1993) and *K. terrigena* (Greiner et al., 1997). It was previously reported that positive monovalent cations did not influence on the alkaline phosphatase activity, while positive bivalent cations had different effects on the enzyme; Mg²⁺, Ca²⁺, Co²⁺, and Mn²⁺ activated the enzyme while Zn²⁺, Cu²⁺, and Cd²⁺ inhibited this enzyme (Kim et al., 2002).

Previous reports showed that SDS showed a strong inhibitory effect on other phytases, such as *A. niger* van Teighem phytase, *Eupenicillium parvum* BCC17694 phytase, and *S. thermophile* phytase (Vats et al., 2005; Singh et al., 2009; Fugthong et al., 2010). For example, *A. niger* van Teighem phytase lost 92% of its activity at 0.1% SDS.

Previous reports showed that metal ions could affect enzyme activity. Since metal ions act as Lewis acids, and coordination complexes of these can have different geometries, they may perform differently toward proteins as a ligand (Dokuzparmak et al., 2017). These differences may also result in metal binding to different sites, and, therefore, change the enzyme structure in different ways and affect the enzyme activity. Neira-Vielma

and co-workers reported that the activity of the *A. niger* phytase increased in the presence of Zn⁺² (Neira-Vielma et al., 2018), while phytase activity of *B. subtilis* CF92 (Hong et al., 2011) and *L. plantarum*, (Sumengen et al., 2012) was also inhibited in the presence of Zn⁺².

Phytases of *K. pneumoniae* 9-3B and *Bacillus subtilis* were stimulated by EDTA (Kerovuo et al., 2000; Choi et al., 2001). It might be owing to the chelating of metal ions from the phytase complex, which decreases the Km. Inhibition of alkaline phytase activity by EDTA has also been reported for the enzymes produced by *Bacillus* sp. DS11, *Bacillus subtilis* (Kim et al., 1998; Vohra and Satyanarayan, 2003), and *Lactobacillus brevis* phytase (Casey et al., 2003). The inhibition by EDTA indicates that some divalent ions are important for complete phytase activity. As mentioned above, EDTA removes divalent metal ions from proteins, thereby modifying their conformations (Ornela et al., 2019).

Finally, LOR10 phytase has engaging activity and stability under acidic pH, excellent thermal stability, and fine ion activity and stability. So, LOR10 phytase can be applied as feed additives, as it could release inorganic phosphate from feed during the digestion process of animals. Currently, phytase has emerged as the world's most extensively used food enzyme. It had been shown that the phytase enzyme has giant latent in food industrial zone for bread processing, isolation of plant proteins, grains wet milling, and cereal bran fractionation.

ACKNOWLEDGEMENT

The authors express their gratitude to the Research Council of the Shahid Bahonar University of Kerman (Iran) for its financial support during this project.

REFERENCES

- Afinah, S., Yazid, A.M., Anis Shobirin, M.H. & Shuhaimi, M. 2010. Phytase: application in food industry. Food Research International Journal 17: 13-21.
- Azadian, F., Badoei-dalfard, A., Namaki-Shoushtari, A. & Hassanshahian, M. 2016. Purification and biochemical properties of a thermostable, haloalkaline cellulase from *Bacillus licheniformis* AMF-07 and its application for hydrolysis of different cellulosic substrates to bioethanol production. Molecular Biology Research Communications 5: 143-155.
- Badoei-Dalfard, A. & Karami, Z. 2013. Screening and isolation of an organic solvent tolerant-protease from *Bacillus* sp. JER02: activity optimization by response surface methodology. Journal of Molecular Catalysis B: Enzymatic 89: 15-23.
- Badoei-Dalfard, A., Parhamfar, M. & Karami, Z. 2019. Characterization of a thermostable,

- acidic-phytase from *Bacillus tequilensis* Dm018; medium optimization by response surface methodology. *Catalysis Letters* 149: 2961-2972.
- Baruah, A., Sahu, N.P., Pal, A.K., Debnath, D., Yengkokpam, S. & Mukherjee, S.C.** 2007. Interactions of dietary microbial phytase, citric acid and crude protein level on mineral utilization by Rohu, *Labeo rohita* (Hamilton), Juveniles. *Journal of the World Aquaculture Society* 38: 238-249.
- Cao, L., Wang, W., Yang, C., Yang, Y., Diana, J. & Yakupitiyage, A.** 2007. Application of microbial phytase in fish feed. *Enzyme and Microbial Technology* 40: 497-507.
- Casey, A. & Walsh, G.** 2004. Identification and characterization of a phytase of potential commercial interest. *Journal of Biotechnology* 110: 313-322.
- Choi, Y.M., Suh, H.J. & Kim, J.M.** 2001. Purification and properties of extracellular phytase from *Bacillus* sp. KHU-10. *Journal of Protein Chemistry* 20: 287-292.
- Chunshan, Q., Linghua, Z., Yunji, W. & Yoshiyuki, O.** 2001. Production of phytase in slow phosphate medium by a novel yeast *Candida krusei*. *Journal of Bioscience and Bioengineering* 92: 154-160.
- Demir, Y., Dikba, N. & Beydemir, Ş.** 2018. Purification and biochemical characterization of phytase enzyme from *Lactobacillus coryniformis* (MH121153). *Molecular Biotechnology* 60: 783-790.
- Dokuzpınar, E., Sirin, Y., Cakmak, U. & Saglam-Ertunga, N.** 2017. Purification and characterization of a novel thermostable phytase from the thermophilic *Geobacillus* sp. TF16. *International Journal of Food Properties* 20: 1104-1116.
- Farahmand, S., Fatemi, F. & Hajhosseini, R.** 2019. Sequencing of the *rus* gene before and after the mutation with DES in the bacterial *Acidithiobacillus* sp. FJ2. *Nova Biologica Reperta* 6: 50-60. (In Persian).
- Farhat, A., Chouayekh, H., Farhat, M.B., Bouchaala, K. & Bejar, S.** 2008. Gene cloning and characterization of a thermostable phytase from *Bacillus subtilis* US417 and assessment of its potential as a feed additive in comparison with a commercial enzyme. *Molecular Biotechnology* 40: 127-135.
- Fugthong, A., Boonyapakron, K., Sornlek, W., Tanapongpipat, S., Eurwilaichitr, L. & Pootanakit, K.** 2010. Biochemical characterization and *in vitro* digestibility assay of *Eupenicillium parvum* (BCC17694) phytase expressed in *Pichia Pastoris*. *Protein Expression and Purification* 70: 60-67.
- Gessler, N.N., Serdyuk, E.G., Isakova, E.P. & Deryabina, Y.I.** 2018. Phytases and the prospects for their application. *Applied Biochemistry and Biotechnology* 54: 352-360.
- Greiner, R., Haller, E., Konietzny, U. & Jany, K.D.** 1997. Purification and characterization of a phytase from *Klebsiella terrigena*. *Archives of Biochemistry and Biophysics* 341: 201-206.
- Greiner, R., Konietzny, U. & Jany, K.D.** 1993. Purification and characterization of two phytases from *Escherichia coli*. *Archives of Biochemistry and Biophysics* 303: 107-113.
- Hong, S.W., Chu, I.H. & Chung, K.S.** 2011. Purification and biochemical characterization of thermostable phytase from newly isolated *Bacillus Subtilis* CF92. *Journal of the Korean Society for Applied Biological Chemistry* 54: 89-94.
- In, M., Jang, J.E.S., Kim, Y.J. & Oh, N.S.** 2004. Purification and properties of an extracellular acid phytase from *Pseudomonas fragi* Y9451. *Journal of Microbiology and Biotechnology* 14: 1004-1008.
- Kerovuo, J., Lappalainen, I. & Reinikainen, T.** 2000. The metal dependence of *Bacillus subtilis* phytase. *Biochemical and Biophysical Research Communications* 268: 365-369.
- Kim Y.O., Kim, H.K., Bae, K.S., Yu, J.H. & Oh, T.K.** 1998. Purification and properties of a thermostable phytase from *Bacillus* sp. DS11. *Enzyme and Microbial Technology* 22: 2-7.
- Kim, Y.H., Gwon, M.N., Yang, S.Y., Park, T.K., Kim, C.G., Kim, C.W. & Song, M.D.** 2002. Isolation of phytase-producing *Pseudomonas* sp. and optimization of its phytase production. *Journal of Microbiology and Biotechnology* 12: 279-285.
- Kim, Y.O., Kim, H.K., Bae, K.S., Yu, J.H. & Oh, T.K.** 1998. Purification and properties of a thermostable phytase from *Bacillus* sp. DS11. *Enzyme and Microbial Technology* 22: 2-7.
- Konietzny, U. & Greiner, R.** 2004. Bacterial phytase: potential application, *in vivo* function and regulation of its synthesis. *Brazilian Journal of Microbiology* 35: 11-18.
- Kumar, V., Sinha, A.K., Makkar, H.P.S. & Becker, K.** 2010. Dietary roles of phytate and phytase in human nutrition: a review. *Food Chemistry* 120: 945-959.
- Lan, G.Q., Abdullah, N., Jalaludin, S. & Ho, Y.W.** 2002. Culture conditions influencing phytase production of *Mitsuokella jalaludinii*, a new bacterial species from the rumen of cattle. *Journal of Applied Microbiology* 93: 668-674.
- Lei, X.G. & Porres, J.M.** 2003. Phytase enzymology, applications, and biotechnology. *Biotechnology Letters* 25: 1787-1794.
- Lei, X.G., Weaver, J.D., Mullaney, E.J., Ullah, A.H. & Azain, M.J.** 2013. Phytase, a new life for an old enzyme. *Annual Review of Animal Biosciences* 1: 283-309.
- Neira-Vielmaa, A.A., Aguilar, C.N., Ilyina, A., Contreras-Esquivel, J.C., Carneiroda-Cunha, M.G., Michelena-Álvarez, G. & Martínez-Hernández, J.L.** 2018. Purification and biochemical characterization of an *Aspergillus niger* phytase produced by solid-state fermentation using triticale residues as substrate. *Biotechnology Reports* 17: 49-54.
- Ornela, P.H. O. & Guimarães, L. H. S.** 2019. Purification and characterization of an alkalistable phytase produced by *Rhizopus microsporus* var. *microsporus* in submerged fermentation. *Process Biochemistry* 81: 70-76.

- Parhamfar, M., Badoei-Dalfard, A., Khaleghi, M. & Hassanshahian, M. 2015. Purification and characterization of an acidic, thermophilic phytase from a newly isolated *Geobacillus stearothermophilus* strain DM12. *Progress in Biological Sciences* 5: 61-73.
- Parhamfar, M., Badoei-Dalfard, A., Parhamfar, M. & Fahimi-Rad, S. 2016. Purification and characterization of an extracellular phosphatase enzyme from *Bacillus* spp. *Journal of Cell and Molecular Research* 8: 90-97.
- Powar, V.K. & Jagannathan, V. 1982. Purification and properties of phytate specific phosphatase from *Bacillus subtilis*. *Journal of Bacteriology* 151: 1102-1108.
- Puppala, K.R., Bhavsar, K., Sonalkar, V., Khire, J.M. & Dharne, M.S. 2019. Characterization of novel acidic and thermostable phytase secreting *Streptomyces* sp. (NCIM 5533) for plant growth promoting characteristics. *Biocatalysis and Agricultural Biotechnology* 18: 101-120.
- Raghavendra, P. & Halami, P.M. 2009. Screening, selection and characterization of phytic acid degrading lactic acid bacteria from chicken intestine. *International Journal of Food Microbiology* 133: 129-134.
- Ramezani-Pour, N., Badoei-Dalfard, A., Namaki-Shoushtari, A. & Karami, Z. 2015. Nitrile-metabolizing potential of *Bacillus cereus* strain FA12; nitrilase production, purification, and characterization. *Biocatalysis and Biotransformation* 33: 156-166.
- Saani, D.M., Lawal, O.T. & Enujiugha, V.N. 2019. Purification and characterization of phytase from *Aspergillus fumigatus* isolated from African giant snail (*Achatina fulica*). *Biocatalysis and Agricultural Biotechnology* 17: 225-232.
- Sajidan, A., Farouk, A., Greiner, R., Jungblut, P., Müller, E.C. & Borriss, R. 2004. Molecular and physiological characterization of a 3-phytase from the *Rhizobacterium Klebsiella pneumoniae* ASR1. *Applied Microbiology and Biotechnology* 65: 110-118.
- Sambrook, J. & Russell, D. 2001. *Molecular cloning: a laboratory manual*, Cold Spring Harbor, New York.
- Sapna, J.J. & Singh, B. 2016. Characteristics and biotechnological applications of bacterial phytases. *Process Biochemistry* 51: 159-169.
- Selle, P.H. & Ravindran, R. 2008. Microbial phytase in poultry nutrition. *Livestock Science* 113: 99-122.
- Shimizu, M. 1992. Purification and characterization of a phytase from *Bacillus subtilis* (natto) N-77. *Bioscience Biotechnology and Biochemistry* 56: 1266-1269.
- Sievers, F., Wilm, A., Dineen, D.G., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., Thompson, J.D. & Higgins, D.G. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology* 7: 539-547.
- Singh, B. & Satyanarayana, T. 2009. Characterization of a HAP-phytase from a thermophilic mould *Sporotrichum Thermophile*. *Bioresource Technology* 100: 2046-2051.
- Singha, B. & Satyanarayana, T. 2011. Phytases from thermophilic molds: their production, characteristics and multifarious applications. *Process Biochemistry* 46: 1391-1398.
- Songa, H.Y., Sheikha, A. F. E. & Hu, D.M. 2019. The positive impacts of microbial phytase on its nutritional applications. *Trends in Food Science and Technology* 86: 553-562.
- Sreedevi, S. & Reddy, B.N. 2012. Isolation, screening and optimization of phytase production from newly isolated *Bacillus* sp.C43. *International Journal of Pharma and Bio Sciences* 2: 218-231.
- Sumengen, M., Dincer, S. & Kaya, A. 2012. Phytase production from *Lactobacillus brevis*. *Turkish Journal of Biology* 36: 533-541.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. 2007. MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* 24: 1596-1599.
- Tye, A.J., Siu, F.K., Leung, T.Y. & Lim, B.L. 2002. Molecular cloning and the biochemical characterization of two novel phytases from *B. subtilis* 168 and *B. licheniformis*. *Applied Microbiology and Biotechnology* 59: 190-197.
- Vasudevan, U.M., Jaiswal, A.K., Krishna, S. & Pandey, A. 2019. Thermostable phytase in feed and fuel industries. *Bioresource Technology* 278: 400-407.
- Vasudevan, U.M., Krishna, S., Jalaja, V. & Pandey, A. 2017. Microbial phytase: impact of advances in genetic engineering in revolutionizing its properties and applications. *Bioresource Technology* 245: 1790-1799.
- Vats, P. & Banerjee, U. 2005. Biochemical characterization of extracellular phytase (myo-inositol hexakisphosphate phosphohydrolase) from a hyper-producing strain of *Aspergillus Niger* van Tieghem. *Journal of Industrial Microbiology and Biotechnology* 32: 141-147.
- Vohra, A. & Satyanarayana, T. 2003. Phytases: microbial sources, production, purification, and potential biotechnological applications. *Critical Reviews in Biotechnology* 23: 29-60.

How to cite this article:

Badoei-Dalfard, A. & Parhamfar, M. 2021. Production and biochemical characterization of a thermostable phytase from *Bacillus amyloliquefaciens* LOR10. *Nova Biologica Reperta* 7: 390-399.